

ADMA, neutrophil to lymphocyte, platelet to lymphocyte ratios and phase angle: effects on inflammation and nutrition in hemodialysis patients

Bahar Gürlek Demirci[✉], Mine Şebnem Karakan[✉]

Department of Nephrology, Ankara Yıldırım Beyazıt University, Faculty of Medicine, Ankara, Türkiye

ABSTRACT

Objectives: Neutrophil/lymphocyte ration (NLR) and platelet/lymphocyte ratio (PLR) levels can be used as systemic inflammatory parameters. Asymmetric dimethyl arginine (ADMA) inhibits endothelial nitric oxide synthase. Phase Angle (PhA) is a potential parameter to screen for inflammatory abnormalities. In present study we aimed to determine the relations between NLR, PLR, ADMA, and PhA in terms of early markers for nutritional status in addition to their well-known role in inflammation.

Methods: A total of 89 patients undergoing maintenance hemodialysis 3 days a week at least 6 months were enrolled. To assess nutritional status, we performed the dietary questionnaire and mini nutritional assessment score (MNAS). ADMA was measured by ELISA. NLR and PLR are calculated from monthly complete blood count tests. Patients were divided into 2 groups according to NLR levels as group 1 (NLR \geq 4.6; n=48) ve and group 2 (NLR<4.6, n=41).

Results: The mean ADMA level was 0.03 ± 0.01 μ mol/L, the mean PhA was $7.2\pm1.1^\circ$. In subgroup analysis, MNAS, albumin levels and phase angle of patients in group 1 were lower and CRP, PLR, ADMA levels were higher when compared to group 2. In correlation analysis, NLR was positively correlated with PLR, CRP and ADMA however negatively correlated with albumin and PhA levels. In regression analysis, NLR, PLR and ADMA were detected as independent predictors of MNAS.

Conclusion: In conclusion our study suggests that NLR, PLR and ADMA are independent predictors for nutritional status and inflammation in patients ongoing hemodialysis.

Keywords: ADMA neutrophil, lymphocyte, platelet, phase angle, inflammation, nutrition, hemodialysis

Maintenance hemodialysis (MHD) is the most applied treatment option for end stage kidney disease [1]. Previous studies have shown that inflammation could be resulted as malnutrition in this population [2]. Several factors as inadequate food intake, uremic toxins, comorbid diseases, the dialysis procedure, inflammatory conditions are all

involved in the development of malnutrition [3]. The prevalence of malnutrition varies up to 76% in MHD patients and considered significantly important for morbidity and mortality of this population [4].

Neutrophil to lymphocyte ratio (NLR) is an inflammatory parameter that shows the presence of systemic inflammation [5]. Previous reports have shown

Corresponding author: Bahar Gürlek Demirci, MD., Assoc. Prof.,
Phone: +90 312 906 25 77, E-mail: bahargurlek@gmail.com

How to cite this article: Gürlek Demirci B, Karakan MŞ. ADMA, neutrophil to lymphocyte, platelet to lymphocyte ratios and phase angle: effects on inflammation and nutrition in hemodialysis patients. Eur Res J. 2024. doi: 10.18621/eurj.1404126

Received: December 12, 2023

Accepted: March 5, 2024

Published Online: May 7, 2024

Copyright © 2024 by Prusa Medical Publishing
Available at <https://dergipark.org.tr/en/pub/eurj>



This is an open access article distributed under the terms of [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/)

that NLR plays a significant role in the development of arteriosclerosis [6, 7]. Besides studies also reported that NLR in MHD patients was related to malnutrition [8]. Platelet to lymphocyte ratio (PLR) is another parameter that is associated with inflammation in patients on MHD [9].

Asymmetric dimethyl arginine (ADMA) is competitive inhibitor of endothelial nitric oxide synthase [10]. Phase angle (PhA) by bioimpedance analysis is an alternative approach for inflammation [11].

In present study we aimed to determine the relations between NLR, PLR, ADMA and PhA in terms of early markers for nutritional status in addition to their well-known role in inflammation.

METHODS

This is a cross-sectional, observational, single-center study aimed to analyze the relationship between inflammation, malnutrition, ADMA, neutrophil to lymphocyte, platelet to lymphocyte ratios and phase angle in MHD patients. Among 130 MHD patients, 89 patients were selected according to following exclusion criteria: (1) lack of regular follow-up, (2) history of chronic inflammatory disease of unknown origin, (3) Kt/V<1.4, (4) active systemic infection or hospitalization in last 3 months, (5) active gastrointestinal disorders, (6) history of peripheral artery disease, (7) active smoking, and (8) history of malignancy. The study was approved by Baskent University Institutional Review Board and Ethics Committee (KA13/245, 12/02/2013). All patients gave informed consent for this study.

Biochemical Assays

All patients' demographical, clinical, and biochemical parameters were analyzed. Venous blood samples were taken after an overnight fast. All blood samples were collected pre dialysis [8]. Lipid profile was measured every 6 months, glycosylated hemoglobin (HbA1C; by high-performance liquid chromatography) levels were measured every 3 months and serum fasting plasma glucose (FPG), uric acid, creatinine, C-reactive protein (CRP), calcium, phosphorus, albumin levels, and complete blood count were measured in monthly periods. NLR and PLR levels were

calculated from monthly complete blood count parameters.

Serum ADMA levels were measured by ELISA method by using an Immunodiagnostic human ADMA kit.

Nutritional Status Assessment

The dietary survey was performed with 3-day dietary histories recorded in a self-completing food diary. The Mini Nutritional Assessment Score (MNAS) includes 18 questions [12].

Phase Angle (PhA) Measurement

After measurement of body weight and height, body composition was measured using the Tanita. Four electrodes were placed on the right hand and foot on the side contralateral to the arteriovenous fistula (if present) of the supine patient. Two electrodes were dorsally placed on the hand: in the metacarpophalangeal articulations and in the corpus, 5cm apart. The pair on the foot were located in the metatarsophalangeal and in the articulation, 6 cm apart. The following were determined: extracellular water (ECW), intracellular water (ICW), and PhA.

Patients were divided into 2 groups according to mean NLR cut-off level as group 1 (NLR \geq 4.6; n=48) and group 2 (NLR<4.6, n=41).

All patients were anuric without any residual urine output which is the most important mortality indicator in this population.

Statistical Analysis

Statistical Package for Social Sciences (version 14.0; SPSS) was used. Kolmogorov-Smirnov test is used for distribution analysis. Normal distributions were expressed as mean (standard deviation). Related data were compared with paired samples t test. Categorical data were compared by χ^2 test, and P<0.05 was considered statistically significant.

RESULTS

This study evaluated 89 patients ongoing MHD. The etiology of CKD in MHD patients was as following: Type 2 DM 38%, hypertension 26%, glomerulonephritis 12%, ADPKD 8 %, and unknown 16 %.

Demographic data of MHD patients were as follows: Patients were 57 (47.8 %) males with mean of age of 58.1 ± 4.8 years. The mean of BMI is 20.4 ± 4.8 kg/m². Duration of dialysis in MHD patients was 12.2 ± 3.6 years. The mean NLR was 4.6 ± 0.7 , the mean PLR was 148.4 ± 28.6 , the mean CRP was 29.4 ± 2.4 mg/L, the mean albumin level was 3.4 ± 0.2 g/dL, the mean ADMA level was 0.4 ± 0.01 μ mol/L, the mean PhA was $7.2 \pm 1.1^\circ$.

According to MNAS of MHD patients, 18.4 % of patients had malnutrition, 42% of patients had risk of malnutrition and 39.6% of patients had normal nutritional status.

In subgroup analysis of study population, demographic data as age, gender, duration of dialysis BMI and ECW/ICW ratio were similar in 2 groups. The MNAS, albumin levels and PhA of patients in group 1 were significantly lower and CRP, PLR, ADMA levels were significantly higher when compared to group 2 (Table 1).

In correlation analysis, NLR was positively correlated with PLR, CRP and ADMA ($r: 0.747$, $P=0.001$, $r: 0.534$, $P=0.003$, and $r: 0.342$, $P=0.001$, respectively), however negatively correlated with albumin and PhA levels ($r: -1.8$, $P=0.016$, $r: -1.4$, $P=0.014$, respectively).

In regression analysis, NLR, PLR and ADMA were detected as independent predictors of MNAS ($P=0.001$, $P=0.01$, and $P=0.02$, respectively).

DISCUSSION

Inflammation plays an essential role in the development and progression of malnutrition in patients ongoing hemodialysis. The peripheral blood cell analysis is preferred for prediction of inflammation and nutritional status [13]. Besides, ADMA is the most potent endogenous inhibitor of nitric oxide synthase and have an important role in inflammation [14]. PhA by is another parameter for inflammatory abnormalities especially in MHD patients.

In present study, we detected ADMA, NLR and PLR are important early markers for nutritional status in addition to their well-known role in inflammation in MHD patients. The incidence of malnutrition in MHD patients reaches up to 60 % in previous studies [15]. Inflammatory parameters as CRP are the most used marker for predicting malnutrition [16]. The negative correlation between inflammation parameters and serum albumin levels are also well known [17]. Recent studies have shown that NLR was also related

Table 1. Subgroup analysis of study population

	Group 1 (n=48)	Group 2 (n=41)	P value
Age (years)	57.4 ± 3.8	59.2 ± 4.4	$0.05 <$
Gender			
Male (n)	31	26	$0.05 <$
Body mass index (kg/m ²)	21.4 ± 2.6	22.1 ± 3.2	$0.05 <$
CRP (mg/dL)	17.8 ± 2.9	5.7 ± 0.7	0.01
Albumin (g/dL)	3.1 ± 0.8	$3. \pm 0.4$	0.03
NLR	4.9 ± 0.2	4.1 ± 0.5	0.01
PLR	154.6 ± 16.6	144.6 ± 21.2	0.03
ADMA (μ m/L)	0.4 ± 0.01	0.28 ± 0.03	0.01
MNAS	$18. \pm 0.4$	25.7 ± 0.4	0.01
ECW/ICW	0.41 ± 0.2	0.43 ± 0.1	$0.05 <$
PhA ($^\circ$)	6.8 ± 1.0	7.7 ± 1.2	0.02

Data are shown as mean \pm standard deviation or number (frequency). ADMA=Asymmetric Dimethylarginine, BMI=Body mass index, CRP=C-Reactive Protein, ECW=Extracellular water, ICW=Intracellular water, NLR=Neutrophil Lymphocyte Ratio, PLR=Platelet Lymphocyte Ratio, MNAS=Mini Nutritional Assessment Score, PhA=Phase Angle

to protein energy wasting. In present study, patients with higher NLR had lower albumin levels and nutritional status and higher inflammation levels as like previous studies. We agree with Malhotra et. al. [18] as leukocytes acts as principal effectors cell in the acute inflammation. Martinez et.al. [19] reported that NLR were significantly associated with nutritional status and inflammation. A recent study suggested that serum albumin was significantly decreased in MHD patients with malnutrition, and NLR was an independent risk factor for malnutrition [20]. In addition, they also demonstrated that NLR was negatively correlated with albumin levels as like present study. They identified these results with the increased NLR might reduce albumin levels, resulting in malnutrition. Our findings are consistent with the previous studies that albumin acts as a negative acute phase parameter in MHD patients.

An elevated PLR level is a novel inflammation marker in chronic kidney disease. In present study, we detected patients with lower nutritional status had higher values of PLR in addition to higher NLR and CRP. A recent study found that patients with poor nutritional status had higher levels of PLR than healthy population [21, 22]. Another study on non-dialysis patients with end stage renal disease detected that either NLR or PLR was positively correlated with CRP. Moreover, they concluded NLR can be an alternative parameter to predict inflammation than PLR in this population [23]. Okayay *et al.* [24] also detected significantly higher NLR in CKD patients than healthy population. Moreover, they showed the presence of higher NLR ratio in both pre-dialysis and dialysis patients, and significantly higher NLR was correlated with higher PLR in CKD patients [24] as like present study. Another study in CKD patients, couldn't detected a statistically significant correlation between SGA score and NLR, PLR, or CRP in patients' group. They showed both ratios of PLR and NLR were positively correlated to high CRP in CKD group when compared to healthy subjects. [25].

ADMA is shown to be a strong predictor of CV disease and mortality in MHD patients [26]. Previous studies reported both malnutrition and inflammation may affect serum ADMA levels especially in chronic kidney disease patients [27]. Moreover, phase angle (PhA) is a simple non-invasive technique that predicts nutritional status [28]. Gener-

ally, a low PhA shows a disrupted ability of store energy [29]. However, PhA may be affected by fluid status especially in MHD patients. In present study ECW/ICW ratios were similar in two groups thus we could regard as PhA as an indicator for nutritional status. In present study, patients with higher NLR and PLR had higher levels of ADMA and lower levels of PhA had worse nutritional status that demonstrates ADMA and PhA may be used as early predictors of nutritional status in MHD patients.

Limitations

The limitations of present study include, firstly, the quite weak statistical power in relation to the small number of patients; secondly, other anthropometric measurements should be added to increase the power of study.

CONCLUSION

In conclusion our study suggests that NLR, PLR and ADMA are independent predictors for nutritional status and inflammation in patients ongoing hemodialysis. Thus, these parameters via complete blood count can be used as an early marker for detecting nutritional status in MHD patients.

Authors' Contribution

Study Conception: BGD; Study Design: BGD; Supervision: MŞK; Funding: BGD; Materials: BGD; Data Collection and/or Processing: BGD; Statistical Analysis and/or Data Interpretation: MŞK; Literature Review: BGD; Manuscript Preparation: BGD and Critical Review: MŞK.

Conflict of interest

The authors disclosed no conflict of interest during the preparation or publication of this manuscript.

Financing

The authors disclosed that they did not receive any grant during conduction or writing of this study.

REFERENCES

1. Kramer A, Pippias M, Noordzij M, et al. The European Renal

- Association-European Dialysis and Transplant Association (ERA-EDTA) Registry Annual Report 2016: a summary. *Clin Kidney J.* 2019;12(5):702-720. doi: 10.1093/ckj/sfz011.
2. Maggio M, Lauretani F. Prevalence, incidence, and clinical impact of cognitive-motoric risk syndrome in Europe, USA, and Japan: facts and numbers update 2019. *J Cachexia Sarcopenia Muscle.* 2019;10(5):953-955. doi: 10.1002/jcsm.12476.
 3. Espahbodi F, Khoddad T, Esmaeili L. Evaluation of malnutrition and its association with biochemical parameters in patients with end stage renal disease undergoing hemodialysis using subjective global assessment. *Nephrourol Mon.* 2014;6(3):e16385. doi: 10.5812/numonthly.16385.
 4. Chertow GM, Johansen KL, Lew N, Lazarus JM, Lowrie EG. Vintage, nutritional status, and survival in hemodialysis patients. *Kidney Int.* 2000;57(3):1176-1181. doi: 10.1046/j.1523-1755.2000.00945.x.
 5. Wang J, Xu MC, Huang LJ, Li B, Yang L, Deng X. Value of neutrophil-to-lymphocyte ratio for diagnosing sarcopenia in patients undergoing maintenance hemodialysis and efficacy of Baduanjin exercise combined with nutritional support. *Front Neurol.* 2023;14:1072986. doi: 10.3389/fneur.2023.1072986.
 6. Pavan A, Calvetti L, Dal Maso A, et al. Peripheral Blood Markers Identify Risk of Immune-Related Toxicity in Advanced Non-Small Cell Lung Cancer Treated with Immune-Checkpoint Inhibitors. *Oncologist.* 2019;24(8):1128-1136. doi: 10.1634/theoncologist.2018-0563.
 7. Bonaventura A, Carbone F, Liberale L, et al. Platelet-to-lymphocyte ratio at the time of carotid endarterectomy is associated with acute coronary syndrome occurrence. *J Cardiovasc Med (Hagerstown).* 2020;21(1):80-82. doi: 10.2459/JCM.0000000000000869.
 8. Chen M, Zheng SH, Yang M, Chen ZH, Li ST. The diagnostic value of preoperative inflammatory markers in craniopharyngioma: a multicenter cohort study. *J Neurooncol.* 2018;138(1):113-122. doi: 10.1007/s11060-018-2776-x.
 9. Maraj M, Kuśnierz-Cabala B, Dumnicka P, et al. Malnutrition, Inflammation, Atherosclerosis Syndrome (MIA) and Diet Recommendations among End-Stage Renal Disease Patients Treated with Maintenance Hemodialysis. *Nutrients.* 2018;10(1):69. doi: 10.3390/nu10010069.
 10. Shafi T, Hostetter TH, Meyer TW, et al. Serum Asymmetric and Symmetric Dimethylarginine and Morbidity and Mortality in Hemodialysis Patients. *Am J Kidney Dis.* 2017;70(1):48-58. doi: 10.1053/j.ajkd.2016.10.033.
 11. da Silva BR, Gonzalez MC, Cereda E, Prado CM. Exploring the potential role of phase angle as a marker of oxidative stress: A narrative review. *Nutrition.* 2022;93:111493. doi: 10.1016/j.nut.2021.111493.
 12. Vellas B, Guigoz Y, Garry PJ, et al. The Mini Nutritional Assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition.* 1999;15(2):116-122. doi: 10.1016/s0899-9007(98)00171-3.
 13. Goswami KK, Ghosh T, Ghosh S, Sarkar M, Bose A, Baral R. Tumor promoting role of anti-tumor macrophages in tumor microenvironment. *Cell Immunol.* 2017;316:1-10. doi: 10.1016/j.cellimm.2017.04.005.
 14. Oliva-Damaso E, Oliva-Damaso N, Rodriguez-Esparragon F, et al. Asymmetric (ADMA) and Symmetric (SDMA) Dimethylarginines in Chronic Kidney Disease: A Clinical Approach. *Int J Mol Sci.* 2019;20(15):3668. doi: 10.3390/ijms20153668.
 15. Matsuzawa R, Yamamoto S, Suzuki Y, et al. The clinical applicability of ultrasound technique for diagnosis of sarcopenia in hemodialysis patients. *Clin Nutr.* 2021;40(3):1161-1167. doi: 10.1016/j.clnu.2020.07.025.
 16. Pedersen BK. Muscles and their myokines. *J Exp Biol.* 2011 Jan 15;214(Pt 2):337-346. doi: 10.1242/jeb.048074.
 17. Kalantar-Zadeh K, Kopple JD, Humphreys MH, Block G. Comparing outcome predictability of markers of malnutrition-inflammation complex syndrome in haemodialysis patients. *Nephrol Dial Transplant.* 2004;19(6):1507-1519. doi: 10.1093/ndt/gfh143.
 18. Malhotra R, Marcelli D, von Gersdorff G, et al. Relationship of Neutrophil-to-Lymphocyte Ratio and Serum Albumin Levels with C-Reactive Protein in Hemodialysis Patients: Results from 2 International Cohort Studies. *Nephron.* 2015;130(4):263-270. doi: 10.1159/000437005.
 19. Diaz-Martinez J, Campa A, Delgado-Enciso I, et al. The relationship of blood neutrophil-to-lymphocyte ratio with nutrition markers and health outcomes in hemodialysis patients. *Int Urol Nephrol.* 2019;51(7):1239-1247. doi: 10.1007/s11255-019-02166-6.
 20. Wang J, Xu MC, Huang LJ, Li B, Yang L, Deng X. Value of neutrophil-to-lymphocyte ratio for diagnosing sarcopenia in patients undergoing maintenance hemodialysis and efficacy of Baduanjin exercise combined with nutritional support. *Front Neurol.* 2023;14:1072986. doi: 10.3389/fneur.2023.1072986.
 21. Liaw FY, Huang CF, Chen WL, et al. Higher Platelet-to-Lymphocyte Ratio Increased the Risk of Sarcopenia in the Community-Dwelling Older Adults. *Sci Rep.* 2017;7(1):16609. doi: 10.1038/s41598-017-16924-y.
 22. Akbas EM, Hamur H, Demirtas L, Bakirci EM, Ozcicek A, Ozcicek F, Kuyruklyildiz U, Turkmen K. Predictors of epicardial adipose tissue in patients with type 2 diabetes mellitus. *Diabetol Metab Syndr.* 2014 May 9;6:55. doi: 10.1186/1758-5996-6-55.
 23. Li P, Xia C, Liu P, et al. Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in evaluation of inflammation in non-dialysis patients with end-stage renal disease (ESRD). *BMC Nephrol.* 2020;21(1):511. doi: 10.1186/s12882-020-02174-0.
 24. Okyay GU, Inal S, Oneç K, et al. Neutrophil to lymphocyte ratio in evaluation of inflammation in patients with chronic kidney disease. *Ren Fail.* 2013;35(1):29-36. doi: 10.3109/0886022X.2012.734429.
 25. Behairy M, Shawky S, Bawady S, El Kezza G, Ahmed F. Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio in Evaluation of Inflammation and Nutritional Status in Chronic Kidney Disease Patients. *Egyptian J Hosp Med.* 2022;89(1):5814-5823. doi: 10.21608/ejhm.2022.266654.
 26. Shafi T, Hostetter TH, Meyer TW, et al. Serum Asymmetric and Symmetric Dimethylarginine and Morbidity and Mortality in Hemodialysis Patients. *Am J Kidney Dis.* 2017;70(1):48-58. doi: 10.1053/j.ajkd.2016.10.033.
 27. Zoccali C, Bode-Böger S, Mallamaci F, et al. Plasma concentration of asymmetrical dimethylarginine and mortality in patients with end-stage renal disease: a prospective study. *Lancet.* 2001;358(9299):2113-2117. doi: 10.1016/s0140-6736(01)07217-8.
 28. Rinaldi S, Gilliland J, O'Connor C, Chesworth B, Madill J.

Is phase angle an appropriate indicator of malnutrition in different disease states? A systematic review. Clin Nutr ESPEN. 2019;29:1-14. doi: 10.1016/j.clnesp.2018.10.010.

29. Norman K, Stobäus N, Pirlich M, Bosy-Westphal A. Bioelec-

trical phase angle and impedance vector analysis--clinical relevance and applicability of impedance parameters. Clin Nutr. 2012;31(6):854-861. doi: 10.1016/j.clnu.2012.05.008.